



Total protein extraction from mice hippocampi and PC12 cells

INSERM u862, Neurocentre Magendie, University of Bordeaux (Bordeaux, France)

CONTEXT

Glucocorticoid hormones are released during the active phase of the circadian cycle and in response to stress. Stress-induced high levels of glucocorticoids have been shown to increase the memory of stress-associated events and to contribute to the development of stress-related pathologies (depression, anxiety, drug abuse...) This study provides a complete molecular pathway (GR/Egr-1/MAPK/Synapsin-Ia-Ib) through which stress and glucocorticoids enhance the memory of stress-related events.


The study involved total protein extractions both from mice hippocampi and PC12 cells using Precellys®24 homogenizer [1].

MATERIAL

- Precellys®24 homogenizer
- Precellys® kit : 03961-1-003 (1.4mm ceramic beads)
- Samples: mice hippocampi and PC12 cells
- Buffer: RIPA buffer containing protease and phosphatase inhibitors (Sigma)

PROTOCOL

- Precellys®24: 5000 rpm, 2x30 sec, 10 sec break
- Centrifugation: 10000 rpm, 10 min, 4°C
- SDS-PAGE, immunoblotting with relevant antibodies.




[1] Revest et al. 2010. The enhancement of stress-related memory by glucocorticoids depends on synapsin-Ia/Ib. *Molecular Psychiatry*, n° 4 (April) doi:10.1038/mp.2010.40



RESULTS

The figure below shows an example of total protein extraction from PC12 cells using Precellys®24.

Total extracts were analyzed by western blot. The corresponding X-ray films were quantified by densitometry.



Corticosterone-induced phosphorylation of synapsin-I depends on Egr-1 activation. Total extracts from PC12 cells were analyzed by western blot 1 and 3 h after treatment with 10nM corticosterone (gray bars). The corresponding X-ray films were quantified by densitometry (optical density, OD, mean ± s.e.m., n = 3). *P < 0.05; ***P < 0.001, in comparison with basal levels.

Those results allowed to better understand the action of corticosterone in the molecular cascade mediating the enhancement of stress-related memories.

CONCLUSION

Precellys® is well adapted for extraction of total proteins from brain samples and cell lines.

Sample homogenization is quick and efficient in term of quantity and quality of extracted proteins (e.g. phosphoproteins). With our protocol, we are able to treat 24 samples in less than 2 minutes by avoiding samples cross contamination.

For more details, please contact precellys@bertin.fr

03712-810-DU035 / July 2010



エムエス機器株式会社

<http://www.technosaurus.co.jp>

- 東京
〒162-0805 東京都新宿区矢来町 113 番地 TEL(03)3235-0661(代) FAX(03)3235-0669
- 大阪
〒532-0005 大阪市淀川区三国本町2丁目12番4号 TEL(06)6396-0501(代) FAX(06)6395-2588
- 福岡
〒812-0054 福岡市東区馬出 1 丁目 2 番 23 号 TEL(092)631-1012(代) FAX(092)641-1285

※会社名および商品名は、各会社の商標または登録商標です。
※本カタログに記載の規格・仕様・外観は予告なく変更する場合がありますので御了承下さい。